



## Research

**Cite this article:** Langwig KE, Frick WF, Hoyt JR, Parise KL, Drees KP, Kunz TH, Foster JT, Kilpatrick AM. 2016 Drivers of variation in species impacts for a multi-host fungal disease of bats. *Phil. Trans. R. Soc. B* **371**: 20150456. <http://dx.doi.org/10.1098/rstb.2015.0456>

Accepted: 7 July 2016

One contribution of 18 to a discussion meeting issue 'Tackling emerging fungal threats to animal health, food security and ecosystem resilience'.

**Subject Areas:**

ecology, health and disease and epidemiology

**Keywords:**

emerging infectious disease, multi-host pathogen, wildlife disease, *Myotis lucifugus*, white-nose syndrome, *Geomyces destructans*

**Author for correspondence:**

Kate E. Langwig  
e-mail: [klangwig@gmail.com](mailto:klangwig@gmail.com)

Electronic supplementary material is available online at <http://dx.doi.org/10.6084/m9.figshare.c.3493050>.

## Drivers of variation in species impacts for a multi-host fungal disease of bats

Kate E. Langwig<sup>1</sup>, Winifred F. Frick<sup>1</sup>, Joseph R. Hoyt<sup>1</sup>, Katy L. Parise<sup>2,3</sup>, Kevin P. Drees<sup>2,3</sup>, Thomas H. Kunz<sup>4</sup>, Jeffrey T. Foster<sup>2,3</sup> and A. Marm Kilpatrick<sup>1</sup>

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of California, EE Biology/EMS, Santa Cruz, CA 95064, USA

<sup>2</sup>Center for Microbial Genetics and Genomics, Northern Arizona University, Flagstaff, AZ 86011-4073, USA

<sup>3</sup>Department of Molecular, Cellular and Biomedical Science, University of New Hampshire, Durham, NH 03824, USA

<sup>4</sup>Department of Biology, Boston University, Boston, MA 02215, USA

**id** KEL, 0000-0001-8318-1238; WFF, 0000-0002-9469-1839; AMK, 0000-0002-3612-5775

Disease can play an important role in structuring species communities because the effects of disease vary among hosts; some species are driven towards extinction, while others suffer relatively little impact. Why disease impacts vary among host species remains poorly understood for most multi-host pathogens, and factors allowing less-susceptible species to persist could be useful in conserving highly affected species. White-nose syndrome (WNS), an emerging fungal disease of bats, has decimated some species while sympatric and closely related species have experienced little effect. We analysed data on infection prevalence, fungal loads and environmental factors to determine how variation in infection among sympatric host species influenced the severity of WNS population impacts. Intense transmission resulted in almost uniformly high prevalence in all species. By contrast, fungal loads varied over 3 orders of magnitude among species, and explained 98% of the variation among species in disease impacts. Fungal loads increased with hibernating roosting temperatures, with bats roosting at warmer temperatures having higher fungal loads and suffering greater WNS impacts. We also found evidence of a threshold fungal load, above which the probability of mortality may increase sharply, and this threshold was similar for multiple species. This study demonstrates how differences in behavioural traits among species—in this case microclimate preferences—that may have been previously adaptive can be deleterious after the introduction of a new pathogen. Management to reduce pathogen loads rather than exposure may be an effective way of reducing disease impact and preventing species extinctions.

This article is part of the themed issue 'Tackling emerging fungal threats to animal health, food security and ecosystem resilience'.

## 1. Introduction

Emerging infectious diseases are an important threat to wildlife populations [1]. Increases in human trade and travel over the last 50 years have driven increases in emerging pathogens, and the introduction of generalist pathogens threaten both human and wildlife populations [2]. Generalist pathogens are capable of infecting multiple host species, which has led to devastation of communities [3,4], species extinctions [5,6] and cascading effects on ecosystems [7,8]. Changes in communities and ecosystems are influenced by variability in population impacts of multi-host pathogens, and some species may even decline to extinction, whereas others suffer little mortality [4,9–12]. Understanding the mechanisms that drive variation in species impacts can help to reduce mortality, prevent species extinctions and thus minimize ecosystem effects [13,14].

The impact of a disease on a population is the product of the fraction of the population infected multiplied by the fraction of infected individuals that die or

fail to reproduce from disease [9,15]. Disease management can target either or both of these components [16]. For humans, public health approaches frequently focus on preventative measures that reduce transmission through behavioural modification or vaccines, whereas medical interventions usually reduce disease severity in infected individuals through drug treatment. Strategies that reduce transmission are preferred if they are more cost-effective than reducing symptoms after exposure. However, in some cases, reducing disease severity by reducing pathogen load can also substantially reduce transmission, particularly for chronic diseases. For example, the discovery that treatment with antiviral drugs greatly reduces infectiousness of HIV patients has revolutionized control of this disease [17].

Long-term control of diseases in wildlife by reducing transmission or disease severity carry additional logistical difficulties in delivering drugs to large enough numbers of free-ranging animals, and challenges in modifying hosts behaviour to reduce transmission. In addition, the relatively short lifespan of many species makes it necessary to have high-frequency repeated interventions that are too costly for all but a few species with a small number of individuals (e.g. vaccination of black-footed ferrets for canine distemper, vaccination of all California condors for West Nile virus [18]). Further, for multi-host pathogens, variation in disease severity is frequently attributed to innate host-specific factors (e.g. immune function), which are not yet easily manipulated for disease management purposes.

However, environmental conditions form the third part of the 'disease triangle', and wildlife disease control may be possible through manipulations that target key aspects of host–pathogen interactions. For example, altering environmental conditions can reduce growth and survival of the pathogen outside the host, and may also increase host defenses [19]. However, identifying environmental factors that can be efficiently targeted requires understanding interactions between hosts, the pathogen and the environment [14]. A detailed understanding of environmental influences on disease may open up a suite of management options that do not require the perpetual actions and funding associated with treatments and vaccination.

Several multi-host fungal pathogens are particularly emblematic of the interplay between host, pathogen and the environment [20–22]. Many fungal pathogens survive outside the host for long periods, or they infect hosts with variable body temperatures and are, therefore, strongly affected by environmental conditions [19,20]. For example, for chytridiomycosis, a fungal disease of amphibians, the highest mortality occurs in cool, high-elevation areas [23]. While few fungal pathogens threaten endothermic immunocompetent species, ectothermic or heterothermic species (such as hibernating mammals, that substantially change core temperatures with seasons) may be more vulnerable to fungal disease. Management of fungal disease has recently been highlighted as a substantial challenge for human, agricultural and wildlife health [20].

White-nose syndrome (WNS), caused by the fungal pathogen *Pseudogymnoascus destructans* [24,25], is a recently emerged fungal disease that has caused widespread mortality in many communities of hibernating bats and is predicted to drive several species extinct [12,26]. WNS was first detected in Schoharie County, New York in 2006, and has since spread across eastern North America (figure 1).

The first detection of *P. destructans* in North America occurred in a commercial tourist cave, and may have been facilitated by human movement [27–29]. *Pseudogymnoascus destructans* is endemic to Eurasia [30–32], where disease severity and population impacts appear to be much lower [33], and several species of Chinese bats have much lower loads than North American species [34]. By the end of the winter of 2015–2016, four bat species distributions within the USA were almost entirely encompassed by the spread of WNS (figure 1). In addition, in March 2016, WNS was detected in Washington state, approximately 1700 km from the nearest known infected site. This spreading event substantially expanded the distribution of the disease (figure 1), and doubled the number of bat species at risk.


Mortality from WNS differs substantially among species, despite them co-occurring at the same sites. Some species declined more than 90% in the first year following WNS detection, whereas population growth rates in other species only decreased 8% [12,35]. Why species suffer such disparate impacts from WNS is a key question that may unlock clues about how to better manage this widespread disease.

Here, we examine how differences in exposure (defined as detection of *P. destructans* on bat skin), fungal loads (the abundance of *P. destructans* on bat skin) and environmental factors determine disease impacts in the six species of bats most impacted by WNS. Previous work has shown that in areas where the disease has been present for multiple years, bats first become infected when they return to hibernacula in the autumn, and both transmission and fungal growth on bats occurs primarily during winter once bats lower their body temperature and begin to hibernate [36]. The relatively high infection prevalence (greater than 50%) observed in many populations of all six species [36] suggests that variation among species in mortality after infection may be especially important in determining population impacts. Although the exact mechanism by which infection with *P. destructans* leads to death is unknown, tissue damage from fungal invasion is thought to set off a cascade of physiological disruptions [37,38], which eventually lead to death approximately 70–120 days after infection [24]. Increases in the extent of tissue invasion with fungal loads are likely to increase pathology and lead to an accelerating probability of death, as has been observed for another fungal pathogen that infects the skin, chytridiomycosis [22]. *Pseudogymnoascus destructans* growth increases with temperature across the range of hibernation temperatures commonly used by bats (approx. 1–12° C) [39–41]. Thus, we hypothesize that species that roost at warmer temperatures will have higher fungal loads and suffer higher mortality and impact from WNS. To test these predictions and hypotheses, we compared patterns of infection prevalence and intensity with differences in species impacts, and then examined links between microclimate temperatures used by bats and fungal loads. Finally, we examined patterns of changes in fungal loads over time to determine whether bats reach a threshold level of fungal infection above which mortality may increase sharply.

## 2. Material and methods

### (a) Field sampling and analysis

We sampled bats at 21 hibernacula in New York, Vermont, Massachusetts, Virginia and Illinois. For five species, we sampled



**Figure 1.** The distribution of white-nose syndrome (hatched) as of March 2016, and distribution maps for six hibernating bat species (colours) in the USA and Canada. Inset: Counts of hibernating bats at sites from 1979 to 2011, with per cent declines in the first year of WNS detection indicated in the upper left of the graph.

bats twice per hibernation season (November/December and March/early April) in sites where *P. destructans* had been detected at least 1 year previously. For Northern long-eared myotis (*Myotis septentrionalis*), we included late hibernation load data for two sites in the first year of WNS detection because exclusion of the invasion year data would result in small sample sizes for this species due to near extirpation in the first year of WNS. We previously used these infection data to examine seasonal changes in prevalence and fungal loads within species [36] (data available from [42]). Here, we examine links between infection patterns and differences in mortality among species [12]. One to six bat species were present in each hibernaculum, including the little brown myotis (*Myotis lucifugus*), Northern long-eared myotis, Eastern small-footed myotis (*Myotis leibii*), Indiana myotis (*Myotis sodalis*), tri-colored bat (*Perimyotis subflavus*) and the big brown bat (*Eptesicus fuscus*) (electronic supplementary material, table S1). At 67% of sites, three of six species co-occurred together, and individuals of different species frequently roosted within 1 m of each other.

We used epidermal swab sampling to determine prevalence and fungal loads of *P. destructans* [36]. Swabs were stored in RNA-later<sup>®</sup> for preservation until extraction. We tested samples for *P. destructans* DNA using real-time polymerase chain reaction (PCR) [43] and quantified fungal loads based on the cycle threshold ( $C_t$ ) value to estimate a fungal load on each bat, with a cut-off of 40 cycles. Quantification of serial dilutions of the DNA from 10 ng to 1000 fg resulted in  $C_t$  scores ranging from 17.33 to 30.74 and a quantification relationship of  $C_t = -3.348 \times \log_{10}(P. destructans \text{ (ng)}) + 22.049$ ,  $r^2 = 0.986$ . We use the term 'infection' to refer to the presence and quantity of *P. destructans* DNA on bat skin, detected using quantitative PCR on epidermal swab samples [20–22,44]. Previous work suggests there is a strong correlation between the abundance of *P. destructans* on bat skin from a swab and tissue invasion as detected by histology [45].

We recorded roost temperature of approximately half of sampled bats (812) using an infrared laser thermometer (Fluke 62 MAX + infrared thermometer) at time of sampling. Measurements were taken of the rock surface directly adjacent

to the roosting bat, while standing within 2 m of the surface. We calibrated on-site roost temperatures with approximately 100 measurements from individual Onset HOBO loggers (model: u23-001; up to 4 per site, twice per year) that recorded a temperature measurement from a single location. Temperature measurements using the infrared laser thermometer were unbiased across a range of temperatures from 1°C to 15°C and within 0.75°C of the HOBO logger temperature.

## (b) Statistical analyses

To estimate prevalence and fungal loads for each species, we used generalized linear mixed models (glmm) with site as a random effect and species interacting with date of sampling as fixed effects (function *glmer* in package *lme4* [46] in R v. 3.02 [47]). We calculated a predicted prevalence and  $\log_{10}$  load for each species on 1 December and 1 March from the glmm model described above to standardize early and late hibernation time points because prevalence and loads increase over winter and bats were sampled at different times [36].

We quantified WNS impact using the change in median population growth rate,  $\Delta\lambda$ , pre-WNS and after the first year of WNS detection, based on previous analyses of 121 colonies of six species of hibernating bats common in the northeastern USA collected over a 30-year period [12]. This dataset provides a large set of sites where count data have been rigorously collected pre- and post-WNS arrival for all six species, enabling us to estimate the change in population growth rate due to disease. These declines are consistent with estimates from other datasets [26,48] including estimates of declines based on summer activity [49,50]. To make comparisons among datasets, we estimated species prevalence and loads, and accounted for site differences by including site as a random effect. We examined the effect of *P. destructans* prevalence and loads on WNS impacts using phylogenetic regression [51] in MATLAB (v. R2013). Species that are closely related often suffer very dissimilar impacts (i.e. little brown and Eastern small-footed bats) [26]. Nonetheless, we accounted for phylogeny in our analyses because the absence of a phylogenetic signal for trees with less than 20 species can be difficult to assess [51]. The approach performs generalized least-squares regression using a variance-covariance matrix expected under an Ornstein–Uhlenbeck model of evolution based on the phylogenetic relationships among species [51]. We used a pruned phylogenetic tree of the six species based on the cytochrome *b* mitochondrial gene [26] (electronic supplementary material, table S2).

We estimated the average roosting temperature for each of the six species using a model of site as a random effect interacting with species (the best fitting model), and averaging the mean predicted temperatures from the regression across all sites. To examine correlations between fungal loads and roosting temperature, we calculated average loads for each species using only data for the individuals for which roosting temperature data were also collected. We then used phylogenetic regression as described above to examine the effect of species roosting temperature on predicted  $\log_{10}$  transformed *P. destructans* loads.

It is worth noting that in both analyses—correlations of WNS impacts and infection prevalence and fungal loads, and roosting temperature and fungal loads—measurement error due to instruments or techniques and the variation among individuals in estimating species' means decreases the power to detect significant relationships, but generally does not lead to an increased probability of a type I statistical error (failing to reject the null hypothesis). As a result, the relationships described below are likely to be even stronger than those reported here. In addition, we performed additional statistical analyses to address this variation, included weighted regression (using the inverse of the

standard error of the estimates) and major axis regression to account for the uncertainty in the estimates of the predictor variables.


Finally, we examined how increases in loads differed among sites and species over winter to determine if fungal loads might reach an asymptote. We estimated the increase in load over winter in individual colonies of each species by fitting a linear model of site interacting with species to the change in *P. destructans* loads. We then compared the change in loads over time for each colony with the early winter loads for that colony. We also examined changes in skewness of load distributions between early and late winter loads for each species.

## 3. Results

We sampled 1314 bats of six species in 21 hibernation sites across New York, Virginia, Massachusetts, Vermont and Illinois (electronic supplementary material, table S1). In early winter, infection prevalence varied from 20% to 90% among species and this was significantly correlated with impacts, measured as the difference in population growth rate before and in the first year after detection of WNS (figure 2a) [12]. Fungal loads on bats at the beginning of hibernation varied by almost 2 orders of magnitude among species and were even more strongly correlated with impacts (figure 2b). By late hibernation, however, prevalence of *P. destructans* had increased to more than 80% for five of six species and was no longer significantly correlated with WNS impacts (figure 2c). By contrast, late hibernation loads of *P. destructans* now varied over 3 orders of magnitude among species and were very strongly correlated with WNS impacts (figure 2d), and neither early nor late prevalence were significant in a model with late hibernation fungal loads (both  $p > 0.45$ ).

Species' fungal loads at the end of hibernation increased significantly with average roosting temperatures (figure 3) and early hibernation prevalence, and there was a marginally significant relationship between hibernation roosting temperature and species impacts ( $0.07 \pm 0.04x + 0.24$ ,  $p = 0.051$ ). Models incorporating both early hibernation prevalence and temperature had slightly higher support than models with either predictor alone ( $\Delta\text{AIC}_C = 0.41$ ). The species with the highest fungal loads, the Northern long-eared bat, roosted at temperatures on average, 6°C warmer than the Eastern small-footed bat (*M. leibii*), which had loads that were 100-fold lower (figure 3). The relationship between roosting temperature and fungal load was even stronger using weighted regression using the inverse of roosting temperature standard error estimates as weights ( $0.26 \pm 0.09x - 4.14$ ,  $p = 0.02$ ), and the results were essentially identical using major axis regression ( $0.26 \pm 0.1x - 4.22$ ;  $p = 0.045$ ).

Average fungal loads within colonies of bats increased during winter, but two lines of evidence suggest that loads on individual bats asymptote, or bats die as they approached loads of approximately  $10^{-1}$  ng. First, the distributions of logged loads shifted from being mostly unskewed (skewness closer to 0) or positive for most species in early hibernation to being more negatively skewed during late hibernation, especially for the three highly impacted species (figure 4; see electronic supplementary material for simulations of how differences in fungal growth on individuals influence fungal load skewness). Loads on Northern long-eared bats at the beginning of winter were already high (approaching the highest recorded fungal loads), and were almost significantly



**Figure 2.** Impact of white-nose syndrome, measured as the change in population growth rates ( $\lambda$ ) following the detection of WNS (based on data and analyses in [12]) and infection prevalence (*a,c*) and intensity (*b,d*) of *P. destructans* for six bat species. Error bars show  $\pm$  s.e. of the predicted mean. Lines show significant relationships, and the dashed line in (*a*) indicates that this relationship was significant only in a univariate regression, but not significant in a model with early or late fungal loads. Equations in the figure report results from phylogenetic regression in the format slope  $\pm$  s.e. of the slope plus the intercept. Late hibernation prevalence was not a significant predictor of WNS impacts (intercept:  $-0.22$ , slope:  $0.92 \pm 0.5$ ,  $p = 0.14$ ,  $R^2 = 0.45$ ).


negatively skewed (bootstrapped confidence intervals,  $p = 0.11$ ). Late winter loads on this species were even more negatively skewed, and few individuals had loads below  $10^{-4}$  ng. Second, the increase in fungal loads for 19 colonies of five species was negatively correlated with mean fungal loads in early hibernation (figure 5). Average loads in colonies that were already high in early hibernation showed little increase over winter, whereas colonies with low average loads increased 1 log or 10-fold each month during hibernation. The best fitting model suggested that the saturation in fungal growth with increasing early hibernation loads was essentially identical for little brown and tri-colored bats (electronic supplementary material, table S3), but increases in fungal loads were lower for Indiana bats.

#### 4. Discussion


The emergence of WNS in North America has altered bat communities on a continental scale, by decimating some species while others have been far less affected [12,26]. Our results suggest that differences in impacts reflect variation in fungal growth resulting in differential pathogen load among species. Differences in timing of infection may initially result in significant differences in exposure and, therefore, infection prevalence at the beginning of hibernation. However, infection prevalence eventually saturates near 100% in most species, which eliminates variation in transmission as a driver of differential population impacts. Instead, different hibernation microclimate preferences

among species appear to result in differential fungal growth on bats over the winter, resulting in wide variation in fungal loads that was tightly correlated with differential population impacts of WNS among bat species.

The relationships between WNS impacts, fungal loads and roosting temperatures offer some insight into the potential mechanisms linking environmental variation and disease impacts. Warmer roosting temperatures, up to  $13^{\circ}\text{C}$ , increase fungal growth rates, with a  $2^{\circ}\text{C}$ – $7^{\circ}\text{C}$  increase in temperature increasing fungal colony size almost fivefold [39]. This provides a potential explanation for the higher fungal loads on bat species roosting at warmer temperatures and, in turn, the higher WNS impacts on these species. Although we cannot infer causation from these relationships, the data are inconsistent with temperature-dependent variation in immune function, which would be predicted to increase with temperature [19]. This would lead to lower fungal loads and impacts at warmer temperatures [52–54]. The causal effect of temperature in increasing WNS impacts among species is also supported by experimental infections of a single species, *M. lucifugus*, at two different temperatures [55]. Furthermore, the wide variation in species impacts within the same genera lends further support that behavioural traits (such as roosting temperature) that exhibit greater individual plasticity [51] may be more important than non-labile traits that are likely to be shared between closely related species. Finally, the energetic expenditure of bats is thought to play an integral role in WNS mortality, because bats suffering from WNS arouse far more frequently than uninfected



**Figure 3.** Estimated *P. destructans* loads in late hibernation and hibernation roosting temperature for six bat species. Species' average load and roosting temperature were restricted to individual bats in which both roost temperatures and load were collected ( $n = 419$ ). Statistics in the figure report results from phylogenetic regression in the format slope  $\pm$  s.e. of the slope plus the intercept.




**Figure 4.** Distributions and skewness of fungal loads on bats in early (solid line) and late hibernation (dashed line). Early and late hibernation skewness differed significantly for little browns ( $p = 0.03$ ), tri-colored ( $p < 0.0001$ ) and big brown bats ( $p = 0.004$ ). Skewness differences were compared by calculating skewness of simulated load distributions from 9999 draws from the grouped winter load data by species, and then comparing the bootstrapped skew differences with our observed skew differences to assess significance.

bats [24,56]. This behaviour can prematurely deplete stored fat, and bats dying of WNS are frequently emaciated [25]. However, energetic models indicate that bats roosting at warmer temperatures (across the range of hibernation temperatures) probably expend less energy during winter because most of the energy expenditure used by bats during hibernation is used during euthermic arousals (70–90%), and thermoregulation during these arousal periods is less energetically costly at warmer temperatures [57]. Nonetheless, relationships between body temperature and torpor patterns are complex and further research is needed to address how changes in roost temperature affect torpor patterns of WNS-affected bats [56].

The increase in WNS impact with roosting temperature suggests that although preferences for higher roosting temperatures may have been beneficial for some species prior to the arrival of WNS, this became maladaptive after *P. destructans* was introduced. It is worth noting that there was considerable variation in roosting temperature within each species (figure 3) and intense mortality from WNS may be selecting for individuals that roost in cooler locations. If preference for roosting temperature is a heritable trait, WNS mortality could drive the evolution of a change in behaviour, as well as select for genes associated with surviving colder winter roosting temperatures. This is one possible explanation for the stabilization of little brown bat populations at some sites in the northeastern United States [12]. Future studies are needed to examine whether temperature and absolute humidity preferences of roosting bats decrease as WNS mortality occurs.

Our results suggest that there is a maximum fungal load above which loads do not increase further, and this maximum value is similar for the two species (little brown and tri-colored bats) that we had sufficient data to compare. One explanation for this pattern is that as loads on individual bats increase, disease-associated pathology and mortality increase sharply as loads approach  $10^{-1}$  ng. A similar phenomenon has been suggested for *Batrachochytrium dendrobatidis*, a fungal pathogen of amphibians [22]. However, it is also possible that the smaller increase in fungal loads in colonies of bats that begin winter at



**Figure 5.** Change in *P. destructans* fungal loads over time plotted against fungal loads at the beginning of hibernation for five species of bats in 19 colonies (where early hibernation prevalence was greater than 50%). Statistical results are shown in electronic supplementary material, table S3.

higher loads is simply due to density-dependent fungal growth on the surface of bats' skin. Quantifying fungal loads over time in an experimental infection study or on marked bats in the field could determine which mechanism is resulting in the threshold we observed in fungal loads.

Understanding the drivers of variation in WNS impacts could be used to more effectively guide management interventions. Our results suggest that a reduction in pathogen loads could reduce disease impacts. Previous work has shown that cooler and drier [12] hibernacula appear to serve as refugia from disease impacts for some populations within two species, possibly because they promote lower fungal loads and thereby increase chances of survival. Although we focused on temperature here, humidity may also contribute to differences in declines among species. If so, manipulating hibernacula entrances to create cooler and drier sites or restricting access to reduce bats' use of the warmer and wetter portions of hibernacula has potential as a single-intervention, long-term solution for management of WNS. By contrast, chemical or biological treatments that reduce loads could be effective in the short term, but if not self-perpetuating, would require continual reapplication making long-term management on a broad-scale challenging [18]. In the past, state managers in New York, Pennsylvania and Wisconsin have manipulated mine entrances to stabilize and alter temperatures (G. Turner 2016, unpublished data; A. Hicks 2007, unpublished data; J. Redell 2015, unpublished data). While there are risks associated with these interventions (i.e. temperatures must be within the physiological limits of bats and unsuitable habit could deter bats), data suggest that a very high percentage of bats roosting at sites with warmer temperatures will perish from WNS [12,26,35,48] and small-scale manipulations could be attempted to test for larger scale feasibility of this as a management tool. While environmental manipulation may not be feasible for all sites it may prove useful for a subset of key sites. Some hibernacula contain large populations (tens of thousands of bats) of multiple species, such that making just a few sites cooler and drier could save diverse populations and help maintain genetic diversity. While WNS has extirpated multiple species from many sites, it has yet to drive any species completely to

extinction, and thus management actions could still conserve the multiple species threatened by WNS.

Differences in disease impacts among species are often assumed to result from inherent differences in disease susceptibility (e.g. cellular pathways or receptors for pathogen binding and replication) [58,59], inherent differences in severity given infection [60–62] or pathogen exposure [12,63–65]. As a result, reductions in transmission often form the basis for the management of wildlife disease [16,18]. Our finding that species impacts were strongly correlated with environmentally driven variation in pathogen growth suggests that efforts to reduce disease impacts by altering environmental factors deserve additional consideration. Disease management via reduction of pathogen loads is a cornerstone of human medical treatments [66], and could be more widely used to manage wildlife diseases like WNS. Our results suggest that environmental manipulation to reduce pathogen growth is an underappreciated tool that could help to prevent disease-caused extinctions in wildlife.

**Ethics.** We followed field hygiene protocols in accordance with United States Fish and Wildlife Service WNS Decontamination Guidelines, and individual state recommendations. All research was conducted under protocol #11-022 approved by the IACUC of Boston University and protocol #Frickw1106 approved by the IACUC of the University of California, Santa Cruz.

**Data accessibility.** Data are available from [42]. For additional data requests contact K. Langwig.

**Authors' contributions.** K.E.L., T.H.K., W.F.F., J.T.F., A.M.K.: conceived and designed the study; K.E.L., J.R.H., A.M.K.: performed the study; K.E.L., J.R.H., K.L.P., K.P.D., A.M.K.: analysed the data; T.H.K., J.T.F.: contributed reagents/materials/analysis tools and K.E.L., A.M.K.: wrote the paper.

**Competing interests.** We declare we have no competing interests.

**Funding.** This work was supported by the National Science Foundation (DEB-1336290, DEB-1115895 and DGE-0741448), Bat Conservation International, the US Fish and Wildlife Service and the National Geographic Society. The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

**Acknowledgements.** We thank R. Reynolds, W. Orndorf, K. Powers (VA), S. Darling (VT), C. Herzog, R. von Linden, K. O'Conner (NY), D. Kirk, J. Kath (IL) and T. Cheng for assistance in data collection.

## References

- Daszak P, Cunningham AA, Hyatt AD. 2000 Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science* **287**, 443–449. (doi:10.1126/science.287.5452.443)
- Woolhouse MEJ, Gowtage-Sequeria S. 2005 Host range and emerging and reemerging pathogens. *Emerg. Infect. Dis.* **11**, 1842–1847. (doi:10.3201/eid1112.050997)
- Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, Turner GG, Butchkoski CM, Kunz TH. 2010 An emerging disease causes regional population collapse of a common North American bat species. *Science* **329**, 679–682. (doi:10.1126/science.1188594)
- Lips KR *et al.* 2006 Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proc. Natl Acad. Sci. USA* **103**, 3165–3170. (doi:10.1073/pnas.0506889103)
- Skerratt LF, Berger L, Speare R, Cashins S, McDonald KR, Phillott AD, Hines HB, Kenyon N. 2007 Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* **4**, 125–134. (doi:10.1007/s10393-007-0093-5)
- Smith KF, Sax DF, Lafferty KD. 2006 Evidence for the role of infectious disease in species extinction and endangerment. *Conserv. Biol.* **20**, 1349–1357. (doi:10.1111/j.1523-1739.2006.00524.x)
- Holdo RM, Sinclair ARE, Dobson AP, Metzger KL, Bolker BM, Ritchie ME, Holt RD. 2009 A disease-mediated trophic cascade in the serengeti and its implications for ecosystem C. *PLoS Biol.* **7**, e1000210. (doi:10.1371/journal.pbio.1000210)
- Whitlaw HA, Lankester MW. 1994 A retrospective evaluation of the effects of parelaprostrongylosis on moose populations. *Can. J. Zool.* **72**, 1–7. (doi:10.1139/z94-001)
- LaDeau SL, Kilpatrick AM, Marra PP. 2007 West Nile virus emergence and large-scale declines of North American bird populations. *Nature* **447**, 710–713. (doi:10.1038/nature05829)
- van Riper C, van Riper SG, Goff ML, Laird M. 1986 The epizootiology and ecological significance of malaria in Hawaiian land birds. *Ecol. Monogr.* **56**, 327–344. (doi:10.2307/1942550)
- Harvell CD *et al.* 1999 Emerging marine diseases—climate links and anthropogenic factors. *Science* **285**, 1505–1510. (doi:10.1126/science.285.5433.1505)
- Langwig KE, Frick WF, Bried JT, Hicks AC, Kunz TH, Marm Kilpatrick A. 2012 Sociality, density-dependence and microclimates determine the persistence of populations suffering from a novel fungal disease, white-nose syndrome. *Ecol. Lett.* **15**, 1050–1057. (doi:10.1111/j.1461-0248.2012.01829.x)
- McCallum H. 2012 Disease and the dynamics of extinction. *Phil. Trans. R. Soc. B* **367**, 2828–2839. (doi:10.1098/rstb.2012.0224)
- Paull SH, Song S, McClure KM, Sackett LC, Kilpatrick AM, Johnson PTJ. 2012 From superspreaders to disease hotspots: linking transmission across hosts and space. *Front. Ecol. Environ.* **10**, 75–82. (doi:10.1890/110111)
- Kilpatrick AM, Peters RJ, Dupuis li AP, Jones MJ, Daszak P, Marra PP, Kramer LD. 2013 Predicted and observed mortality from vector-borne disease in wildlife: West Nile virus and small songbirds. *Biol. Conserv.* **165**, 79–85. (doi:10.1016/j.biocon.2013.05.015)
- Wobeser G. 2002 Disease management strategies for wildlife. *Rev. Sci. Tech. Off. Int. Epizoot.* **21**, 159–178. (doi:10.20506/rst.21.1.1326)
- Cohen MS *et al.* 2011 Prevention of HIV-1 infection with early antiretroviral therapy. *N. Engl. J. Med.* **365**, 493–505. (doi:10.1056/NEJMoa1105243)
- Langwig KE *et al.* 2015 Context dependent conservation responses to wildlife disease. *Front. Ecol. Environ.* **13**, 195–202. (doi:10.1890/140241)
- Raffel TR, Romansic JM, Halstead NT, McMahon TA, Venesky MD, Rohr JR. 2012 Disease and thermal acclimation in a more variable and unpredictable climate. *Nat. Clim. Change* **3**, 146–151. (doi:10.1038/nclimate1659)
- Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. 2012 Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**, 186–194. (doi:10.1038/nature10947)
- Briggs CJ, Knapp RA, Vredenburg VT. 2010 Zoozotic and epizootic dynamics of the chytrid fungal pathogen of amphibians. *Proc. Natl Acad. Sci. USA* **107**, 9695–9700. (doi:10.1073/pnas.0912886107)
- Vredenburg VT, Knapp RA, Tunstall TS, Briggs CJ. 2010 Dynamics of an emerging disease drive large-scale amphibian population extinctions. *Proc. Natl Acad. Sci. USA* **107**, 9689–9694. (doi:10.1073/pnas.0914111107)
- Kilpatrick AM, Briggs CJ, Daszak P. 2010 The ecology and impact of chytridiomycosis: an emerging disease of amphibians. *Trends Ecol. Evol.* **25**, 109–118. (doi:10.1016/j.tree.2009.07.011)
- Warnecke L, Turner JM, Bollinger TK, Lorch JM, Misra V, Cryan PM, Wibbelt G, Blehert DS, Willis CKR. 2012 Inoculation of bats with European *Geomyces destructans* supports the novel pathogen hypothesis for the origin of white-nose syndrome. *Proc. Natl Acad. Sci. USA* **109**, 6999–7003. (doi:10.1073/pnas.1200374109)
- Lorch JM *et al.* 2011 Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature* **480**, 376–378. (doi:10.1038/nature10590)
- Frick WF *et al.* 2015 Disease alters macroecological patterns of North American bats. *Glob. Ecol. Biogeogr.* **24**, 741–749. (doi:10.1111/geb.12290)
- Blehert DS *et al.* 2009 Bat white-nose syndrome: an emerging fungal pathogen? *Science* **323**, 227–227. (doi:10.1126/science.1163874)
- Leopardi S, Blake D, Puechmaille SJ. 2015 White-nose syndrome fungus introduced from Europe to North America. *Curr. Biol.* **25**, R217–R219. (doi:10.1016/j.cub.2015.01.047)
- Puechmaille SJ, Frick WF, Kunz TH, Racey PA, Voigt CC, Wibbelt G, Teeling EC. 2011 White-nose syndrome: is this emerging disease a threat to European bats? *Trends Ecol. Evol.* **26**, 570–576. (doi:10.1016/j.tree.2011.06.013)
- Hoyt JR *et al.* 2015 Widespread bat white-nose syndrome fungus, northeastern China. *Emerg. Infect. Dis.* **22**, 140–142. (doi:10.3201/eid2201.151314)
- Puechmaille SJ, Verdeyroux P, Fuller H, Gouilh MA, Bekaert M, Teeling EC. 2010 White-nose syndrome fungus (*Geomyces destructans*) in bat, France. *Emerg. Infect. Dis.* **16**, 290–293. (doi:10.3201/eid1602.091391)
- Zukal J *et al.* 2014 White-nose syndrome fungus: a generalist pathogen of hibernating bats. *PLoS ONE* **9**, e97224. (doi:10.1371/journal.pone.0097224)
- Puechmaille SJ *et al.* 2011 Pan-European distribution of white-nose syndrome fungus (*Geomyces destructans*) not associated with mass mortality. *PLoS ONE* **6**, e19167. (doi:10.1371/journal.pone.0019167)
- Hoyt JR *et al.* 2016 Host persistence or extinction from emerging infectious disease: insights from white-nose syndrome in endemic and invading regions. *Proc. R. Soc. B* **283**, 20152861. (doi:10.1098/rspb.2015.2861)
- Langwig KE, Hoyt JR, Parise KL, Kath J, Kirk D, Frick WF, Foster JT, Kilpatrick AM. 2015 Invasion dynamics of white-nose syndrome white-nose syndrome fungus, midwestern United States, 2012–2014. *Emerg. Infect. Dis.* **21**, 1023–1026. (doi:10.3201/eid2106.150123)
- Langwig KE *et al.* 2015 Host and pathogen ecology drive the seasonal dynamics of a fungal disease, white-nose syndrome. *Proc. R. Soc. B* **282**, 20142335. (doi:10.1098/rspb.2014.2335)
- Warnecke L, Turner JM, Bollinger TK, Misra V, Cryan PM, Blehert DS, Wibbelt G, Willis CKR. 2013 Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing damage to mortality. *Biol. Lett.* **9**, 20130177. (doi:10.1098/rsbl.2013.0177)
- Verant ML, Carol MU, Speakman JR, Cryan PM, Lorch JM, Blehert DS. 2014 White-nose syndrome initiates a cascade of physiologic disturbances in the hibernating bat host. *BMC Physiol.* **14**, 227. (doi:10.1186/s12899-014-0010-4)
- Verant ML, Boyles JG, Waldrep W, Wibbelt G, Blehert DS. 2012 Temperature-dependent growth of *Geomyces destructans*, the fungus that causes bat white-nose syndrome. *PLoS ONE* **7**, e46280. (doi:10.1371/journal.pone.0046280)
- Brack V. 2007 Temperatures and locations used by hibernating bats, including *Myotis sodalis* (Indiana bat), in a limestone mine: implications for conservation and management. *Environ. Manage.* **40**, 739–746. (doi:10.1007/s00267-006-0274-y)



41. Webb PJ, Speakman JR, Racey PA. 1996 How hot is a hibernaculum? A review of the temperatures at which bats hibernate. *Can. J. Zool. Rev. Can. Zool.* **74**, 761–765. (doi:10.1139/z96-087)
42. Langwig KE, Frick WF, Hoyt JR, Parise KL, Drees KP, Kunz TH, Foster JT, Kilpatrick AM. 2016 Drivers of variation in species impacts for a multi-host fungal disease of bats. *Dryad Digital Repository* (doi:10.1098/rstb.2015.0456)
43. Muller LK, Lorch JM, Lindner DL, O'Connor M, Gargas A, Blehert DS. 2013 Bat white-nose syndrome: a real-time TaqMan polymerase chain reaction test targeting the intergenic spacer region of *Geomyces destructans*. *Mycologia* **105**, 253–259. (doi:10.3852/12-242)
44. Haas CN, Rose JB, Gerba CP. 1999 *Quantitative microbial risk assessment*, p. 449, 1st edn. New York, NY: John Wiley & Sons, Inc.
45. McGuire LP *et al.* 2016 White-nose syndrome disease severity and a comparison of diagnostic methods. *EcoHealth* **13**, 60–71. (doi:10.1007/s10393-016-1107-y)
46. Bates D, Maechler M, Bolker B. 2011 lme4: Linear mixed-effects models using Eigen and Eigen. R package version 0.999375-42. See <http://CRAN.R-project.org/package=lme4>.
47. R Development Core Team. 2012 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
48. Turner GG, Reeder DM, Coleman JTH. 2011 A five-year assessment of mortality and geographic spread of white-nose syndrome in North American bats and look to the future. *Bat Res. News* **52**, 13–27.
49. Dzal Y, McGuire LP, Veselka N, Fenton MB. 2011 Going, going, gone: the impact of white-nose syndrome on the summer activity of the little brown bat (*Myotis lucifugus*). *Biol. Lett.* **7**, 392–394. (doi:10.1098/rsbl.2010.0859)
50. Brooks RT. 2011 Declines in summer bat activity in central New England 4 years following the initial detection of white-nose syndrome. *Biodivers. Conserv.* **20**, 2537–2541. (doi:10.1007/s10531-011-9996-0)
51. Blomberg SP, Garland T, Ives AR. 2003 Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* **57**, 717–745. (doi:10.1111/j.0014-3820.2003.tb00285.x)
52. Moore MS, Reichard JD, Murtha TD, Nabhan ML, Pian RE, Ferreira JS, Kunz TH. 2013 Hibernating little brown myotis (*Myotis lucifugus*) show variable immunological responses to white-nose syndrome. *PLoS ONE* **8**, e58976. (doi:10.1371/journal.pone.0058976)
53. Bouma HR, Carey HV, Kroese GM. 2010 Hibernation: the immune system at rest? *J. Leukoc. Biol.* **88**, 619–624. (doi:10.1189/jlb.0310174)
54. Bouma HR *et al.* 2011 Low body temperature governs the decline of circulating lymphocytes during hibernation through sphingosine-1-phosphate. *Proc. Natl Acad. Sci. USA* **108**, 2052–2057. (doi:10.1073/pnas.1008823108)
55. Johnson SA, Brack V, Rolley RE. 1998 Overwinter weight loss of Indiana bats (*Myotis sodalis*) from hibernacula subject to human visitation. *Am. Midland Nat.* **139**, 255–261. (doi:10.1674/0003-0031(1998)139[0255:owloib]2.0.co;2)
56. Reeder DM *et al.* 2012 Frequent arousal from hibernation linked to severity of infection and mortality in bats with white-nose syndrome. *PLoS ONE* **7**, e38920. (doi:10.1371/journal.pone.0038920)
57. Jonasson KA, Willis CKR. 2012 Hibernation energetics of free-ranging little brown bats. *J. Exp. Biol.* **215**, 2141–2149. (doi:10.1242/jeb.066514)
58. Pepin KM, Lass S, Pulliam JR, Read AF, Lloyd-Smith JO. 2010 Identifying genetic markers of adaptation for surveillance of viral host jumps. *Nat. Rev. Microbiol.* **8**, 802–813. (doi:10.1038/nrmicro2440)
59. Pulliam JR, Dushoff J. 2009 Ability to replicate in the cytoplasm predicts zoonotic transmission of livestock viruses. *J. Infect. Dis.* **199**, 565–568. (doi:10.1086/596510)
60. Fraser C, Hollingsworth TD, Chapman R, de Wolf F, Hanage WP. 2007 Variation in HIV-1 set-point viral load: epidemiological analysis and an evolutionary hypothesis. *Proc. Natl Acad. Sci. USA* **104**, 17 441–17 446. (doi:10.1073/pnas.0708559104)
61. Komar N, Langevin S, Hinten S, Nemeth NM, Edwards E, Hettler DL, Davis BS, Bowen RA, Bunning ML. 2003 Experimental infection of North American birds with the New York 1999 strain of West Nile virus. *Emerg. Infect. Dis.* **9**, 311–322. (doi:10.3201/eid0903.020628)
62. Yorinks N, Atkinson CT. 2000 Effects of malaria on activity budgets of experimentally infected juvenile Apapane (*Himatione sanguinea*). *Auk* **117**, 731–738. (doi:10.1642/0004-8038(2000)117[0731:EOMOAB]2.0.CO;2)
63. Altizer S, Hochachka WM, Dhondt AA. 2004 Seasonal dynamics of mycoplasmal conjunctivitis in eastern North American house finches. *J. Anim. Ecol.* **73**, 309–322. (doi:10.1111/j.0021-8790.2004.00807.x)
64. Johnson PTJ, Hartson RB, Larson DJ, Sutherland DR. 2008 Diversity and disease: community structure drives parasite transmission and host fitness. *Ecol. Lett.* **11**, 1017–1026. (doi:10.1111/j.1461-0248.2008.01212.x)
65. Wendland LD *et al.* 2010 Social behavior drives the dynamics of respiratory disease in threatened tortoises. *Ecology* **91**, 1257–1262. (doi:10.1890/09-1414.1)
66. Read AF, Day T, Huijben S. 2011 The evolution of drug resistance and the curious orthodoxy of aggressive chemotherapy. *Proc. Natl Acad. Sci. USA* **108**, 10 871–10 877. (doi:10.1073/pnas.1100299108)